



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DESCRIPTIONS OF TWO NEW FORMS OF PERIDERIS.

BY HENRY A. PILSBRY.

Perideris Kobelti n. sp.

Shell ovate, ventricose, whitish under an olivaceous yellow cuticle, brownish in places. Surface obliquely plicatulate in the direction of growth lines, and above the periphery decussated by numerous rather inconspicuous spiral impressed lines. Spire abruptly contracted above, mucronate. Whorls about $6\frac{1}{2}$, the earlier $2\frac{1}{2}$ forming a mamillar projection, the rest rapidly widening, last whorl swollen. Suture white from loss of cuticle, and minutely beaded. Aperture oblique, reddish inside; lip obtuse, brown; columella whitish, vertical rather straight, subtruncate below. Alt. 65, diam. 40, alt. of aperture 35 mm.

Cape Palmas, West Africa. Type in coll. A. N. S. P.

This species is evidently identical with Kobelt's *P. sauleydi* (Conchyl. Cab., p. 42, pl. 13, f. 1, 2). It is far from being the species so called by Joannis. It is likely that Kobelt had not seen Joannis' original description and figure, from the fact that his reference to that author is altogether incorrect; and thus the misidentification arose.

Perideris Sauleydi var. **normalis** n. var.

Shell resembling *Achatina Sauleydi* Joannis (Magazin de Zoologie, 1834, Classe V, pl. 50) in the general coloration, being white streaked with livid purple, with some bluish suffusion, the penultimate whorl whitish with reddish flames. Whorls about $7\frac{1}{2}$, the earlier 3 forming a mamillar mucro, higher and more distinctly differentiated than in *Sauleydi*, those following forming a more tapering cone than in *Sauleydi*, the last not perceptibly constricted below the suture. Suture margined below by a narrow crenate or beaded band. Sculpture: unequally, obliquely plicatulate, sometimes with subobsolete spirals on penultimate whorl, the last whorl with faint, obliquely descending, scar-like impressions at right angles to the growth-lines. Aperture oblique, dark purplish-brown inside; parietal wall orange-brown from the retention of the cuticle by the parie-

tal glaze. Columella somewhat concave, passing gradually into the thin basal lip.

Alt. 76, diam. 40, alt. of aperture, 36 mm.

Alt. 74, diam. 40, alt. of aperture, 35 mm.

"Taboo, Africa" (Robert Swift coll. in A. N. S. Phila.).

This form differs from *P. Sauleydi* (Joannis) in being more elongated and slender throughout, the last whorl not concave above, aperture consequently not acuminate posteriorly as in that species; the spire more elongated and tapering, and the terminal "*mamelon*" more pronounced.

It is also dextral; but as the apparent sinistrality of Joannis' species may possibly (though not probably) be due to an artist's failure to reverse, I do not place great stress upon this feature.

Illustrations of this and the preceding form will appear in the *Manual of Conchology* in due time.